

2014 Recommended Immunizations

for Children from Birth through 6 Years Old

| Birth | 1 month | 2 months | 4 months | 6 months | 12 months | 15 months | 18 months | 19-23 months | 2-3 years | 4-6 years |
|-------|---------|----------|----------|----------|---------------------|-----------|-----------|--------------|-----------|-----------|
| HepB | HepB | | | | HepB | | | | | |
| | | RV | RV | RV | | | | | | |
| | | DTaP | DTaP | DTaP | | DTaP | | | | DTaP |
| | | Hib | Hib | Hib | Hib | | | | | |
| | | PCV | PCV | PCV | PCV | | | | | |
| | | IPV | IPV | IPV | | | | | | IPV |
| | | | | | Influenza (yearly)* | | | | | |
| | | | | | MMR | | | | | MMR |
| | | | | | Varicella | | | | | Varicella |
| | | | | | | HepA** | | | | |

Areas shaded in blue indicate the vaccine can be given during shown age range.

If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.



Is your family growing?
To protect your new baby and yourself against whooping cough, get a Tdap vaccination in the third trimester of each pregnancy. See your doctor for more details.

* Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

** Two doses of Hep A vaccine are needed for lasting protection. The first should be given between 12 and 23 months of age, and the second 6-18 month later.